

A Characterisation of Strong Integer Additive Set-Indexers of Graphs

N K Sudev^{*} and K A Germina[†]

Abstract

An integer additive set-indexer is defined as an injective function $f : V(G) \rightarrow 2^{\mathbb{N}_0}$ such that the induced function $g_f : E(G) \rightarrow 2^{\mathbb{N}_0}$ defined by $g_f(uv) = f(u) + f(v)$ is also injective, where $f(u) + f(v)$ is the sumset of $f(u)$ and $f(v)$. If $g_f(uv) = k \ \forall \ uv \in E(G)$, then f is said to be a k -uniform integer additive set-indexers. An integer additive set-indexer f is said to be a strong integer additive set-indexer if $|g_f(uv)| = |f(u)| \cdot |f(v)| \ \forall \ uv \in E(G)$. We already have some characteristics of the graphs which admit strong integer additive set-indexers. In this paper, we study the characteristics of certain graph classes, graph operations and graph products that admit strong integer additive set-indexers.

Key words: Set-indexers, integer additive set-indexers, strong integer additive set-indexers, Difference sets, nourishing number of a graph.

AMS Subject Classification No.: 05C78

1 Introduction

For all terms and definitions, not defined specifically in this paper, we refer to [7]. Unless mentioned otherwise, all graphs considered here are simple, finite and have no isolated vertices.

Let $\mathbb{N}_0$ denote the set of all non-negative integers. For all $A, B \subseteq \mathbb{N}_0$, the sum of these sets is denoted by $A + B$ and is defined by $A + B = \{a + b : a \in$

^{*}Department of Mathematics, Vidya Academy of Science & Technology, Thalakkottukara, Thrissur - 680501, email: sudev nk@gmail.com

[†]Department of Mathematics, School of Mathematical & Physical Sciences, Central University of Kerala, Kasaragod, email: srgerminaka@gmail.com

$A, b \in B\}$. The set $A + B$ is called the *sumset* of the sets A and B . Also, we have $2A = A + A$.

If either A or B is countably infinite, then their sumset is also countably infinite. Hence, the sets we consider here are all finite sets of non-negative integers. The cardinality of a set A is denoted by $|A|$.

Definition 1.1. [4] An *integer additive set-indexer* (IASI, in short) is defined as an injective function $f : V(G) \rightarrow 2^{\mathbb{N}_0}$ such that the induced function $g_f : E(G) \rightarrow 2^{\mathbb{N}_0}$ defined by $g_f(uv) = f(u) + f(v)$ is also injective.

Definition 1.2. [9] If a graph G has a set-indexer f such that $|g_f(uv)| = |f(u) + f(v)| = |f(u)| \cdot |f(v)|$ for all vertices u and v of G , then f is said to be a *strong IASI* of G .

Definition 1.3. [9] If G is a graph which admits a k -uniform IASI and $V(G)$ is l -uniformly set-indexed, then G is said to have a (k, l) -*completely uniform IASI* or simply a *completely uniform IASI*.

We use the notation $A < B$ in the sense that $A \cap B = \emptyset$. We notice that the relation $<$ is symmetric, but need not be reflexive and transitive. By the sequence $A_1 < A_2 < A_3 < \dots < A_n$, we mean that the given sets are pairwise disjoint.

Lemma 1.4. [9] Let A, B be two non-empty subsets of $\mathbb{N}_0$. Then, $|A + B| = |A| \cdot |B|$ if and only if their difference sets, denoted by D_A and D_B respectively, follow the relation $D_A < D_B$.

Theorem 1.5. [9] Let each vertex v_i of the complete graph K_n be labeled by the set $A_i \in 2^{\mathbb{N}_0}$. Then K_n admits a strong IASI if and only if there exists a finite sequence of sets $D_1 < D_2 < D_3 < \dots < D_n$ where each D_i is the set of all differences between any two elements of the set A_i .

Theorem 1.6. [9] If a graph G admits a strong IASI then its subgraphs also admit strong IASI.

Corollary 1.7. [9] A connected graph G (on n vertices) admits a strong IASI if and only if each vertex v_i of G is labeled by a set A_i in $2^{\mathbb{N}_0}$ and there exists a finite sequence of sets $D_1 < D_2 < D_3 < \dots < D_m$, where $m \leq n$ is a positive integer and each D_i is the set of all differences between any two elements of the set A_i .

2 New Results on Strong IASI Graphs

2.1 Strong IASI of Certain Graph Classes

Based on the results in the above section, we introduce the following notions.

The set D_i of all differences between two elements of the set A_i is called the *difference set* of A_i and the relation $<$ is called the *difference relation* on G . We call the sequence of difference sets, mentioned in Corollary 1.7 a *chain* of difference sets. The *length* of a chain is the number of difference sets in that chain.

For any strong IASI graph with m vertices and n edges, there are m difference sets, one each for each vertex and n relations $<$, one each corresponding to each edge of G . Since G is a strong IASI graph, if two vertices are adjacent in G , then there exists the difference relation between the difference sets of the set-labels of those vertices. These relations forms one or more chain of difference sets. The length of a chain of difference sets is noteworthy and hence we make the following notion.

Definition 2.1. The *nourishing number* of a set-labeled graph is the minimum length of the maximal chain of difference sets in G . The nourishing number of a graph G is denoted by $\varkappa(G)$.

The study about nourishing number of different graphs and graph classes arises much interest. In this section, we discuss about the nourishing number of various graphs.

By Theorem 1.5, we have the following result.

Theorem 2.2. *The nourishing number of a complete graph is n . That is, $\varkappa(K_n) = n$.*

We proceed with the following theorem on the nourishing number of bipartite graphs.

Theorem 2.3. *The nourishing number of a bipartite graph is 2.*

Proof. Let G be a bipartite graph with bipartition (X, Y) which admits a strong IASI. Let $X = \{u_1, u_2, u_3, \dots, u_m\}$ and $Y = \{v_1, v_2, v_3, \dots, v_n\}$. Define $f : V(G) \rightarrow 2^{\mathbb{N}_0}$ such that for two adjacent vertices $u_i \in X$ and $v_j \in Y$, the difference sets of $f(u_i)$ and $f(v_j)$, denoted by D_{u_i} and D_{v_j} respectively, hold the relation $D_{u_i} < D_{v_j}$. Hence, corresponding to each edge in G , there exists a difference relation between the set-labels of its end vertices. Hence, f is a strong IASI and the minimum length of the maximal chain in G is 2. That is, $\varkappa(G) = 2$. $\square$

Proposition 2.4. *The nourishing number of a triangle-free graph is 2.*

Proof. Since G is a triangle-free graph, G can not be a complete graph. Since the relation $<$ is not transitive, we can not find a chain consisting of three or more difference sets corresponding to the set-labels of vertices of G . Hence $\varkappa(G) = 2$. $\square$

A clique of graph G is a complete subgraph of G . We recall that the clique number $\omega(G)$ of a graph is the number of vertices in a maximal clique in G .

Proposition 2.5. *The nourishing number of a graph G is the clique number $\omega(G)$ of G .*

Proof. Let H be the maximal clique of a given graph G . Then, H is complete graph. Hence, by Theorem 1.5, there exists a chain of difference sets $D_1 < D_2 < D_3 < \dots < D_r$, where $r = |V(H)|$. Moreover, no cliques in $G - H$ can have more vertices in G than H . Therefore, $\varkappa(G) = r = \omega(G)$. $\square$

From 2.4 and 2.5, we propose the following result.

Theorem 2.6. *For any strong IASI graph G , the nourishing number $\varkappa(G)$ is given by $\varkappa(G) = |V(H)| = \omega(G)$, where H is the maximal clique of G .*

Interesting questions that arise in this context are about the nourishing number of different graph operations. In the following discussions, we address these problems. First, we check the admissibility of strong IASIs by the union of two graphs and its nourishing number in the following results.

2.2 Strong IASIs of Graph Operations

By the term *last vertex* of a subgraph H of G , we mean a vertex $v \in V(H)$ whose adjacent vertices in G are not in $V(H)$.

Theorem 2.7. *The union $G_1 \cup G_2$ of two graphs G_1 and G_2 , admits a strong IASI if and only if both G_1 and G_2 admit strong IASIs.*

Proof. Let G_1 and G_2 admit strong IASIs say f_1 and f_2 . If G_1 and G_2 are disjoint graphs, then we observe that the IASI f defined by $f(v) = f_i(v)$ if $v \in V(G_i)$, $i = 1, 2$ is a strong IASI for $G_1 \cup G_2$.

If G_1 and G_2 are not disjoint graphs, then re-label the vertices in $G_1 \cap G_2$ in such a way that the difference sets of the set-labels of adjacent vertices in $G_1 \cap G_2$ and the last vertices of $G_1 \cap G_2$ and their adjacent vertices in

G_1 and G_2 follow the relation ' $<$ '. Hence, f is a strong IASI of $G_1 \cup G_2$. Conversely, assume that $G_1 \cup G_2$ admits a strong IASI. Hence, by Theorem 1.6, being the subgraphs of a strong IASI graph $G_1 \cup G_2$, G_1 and G_2 admit the (induced) strong IASIs, $f|_{G_1}$ and $f|_{G_2}$, where $f|_{G_i}$ is the restriction of f to the (sub)graph G_i . $\square$

Theorem 2.8. *Let G_1 and G_2 be two strong IASI graphs. Then, $\kappa(G_1 \cup G_2) \geq \max\{\kappa(G_1), \kappa(G_2)\}$.*

Proof. Let G_1 and G_2 be two strong IASI graphs. Let H_1 and H_2 be the maximal cliques in G_1 and G_2 respectively. If G_1 and G_2 are disjoint, so are H_1 and H_2 . Without loss of generality, let $|V(H_1)| \geq |V(H_2)|$. Then, H_1 is the maximal clique in $G_1 \cup G_2$. Hence, $\kappa(G_1 \cup G_2) = \kappa(G_1)$. Therefore, in general, for disjoint graphs G_1 and G_2 , $\kappa(G_1 \cup G_2) = \max\{\kappa(G_1), \kappa(G_2)\}$. If G_1 and G_2 are not disjoint, then there may exist a subgraph H'_1 , not necessarily complete, in G_1 and a subgraph H'_2 , not necessarily complete, in G_2 such that $H'_1 \cup H'_2 = K_l$, where $l \geq |V(H_1)|, |V(H_2)|$. In this case, $\kappa(G_1 \cup G_2) \geq \max\{\kappa(G_1), \kappa(G_2)\}$. This completes the proof. $\square$

From Theorem 2.8, we observe the following theorem.

Theorem 2.9. *Let G_1 and G_2 be two strong IASI graphs. Then, $\kappa(G_1 \cup G_2) = \max\{\kappa(G_1), \kappa(G_2)\}$ if $G_1 \cap G_2$ is triangle-free.*

Proof. Every complete graph K_n with more than two vertices contains triangles. Hence, since, $G_1 \cap G_2$ is triangle-free, it does not contain any clique. Therefore, there does not exist a subgraph H'_i in $G_i, i = 1, 2$ such that $H'_1 \cup H'_2 = K_l, l > 2$. Hence, $\kappa(G_1 \cup G_2) = \max\{\kappa(G_1), \kappa(G_2)\}$. $\square$

The following theorems check the admissibility of strong IASI by the join of two strong IASI graphs and its nourishing number.

Theorem 2.10. *Let $G_1(V_1, E_1)$ and $G_2(V_2, E_2)$ be two strong IASI graphs. Then, their join $G_1 + G_2$ admits a strong IASI if and only if the difference set of the set-label of every vertex in G_1 is disjoint from the difference sets of the set-labels of all vertices of G_2 .*

Proof. Let $V(G_1) = \{u_1, u_2, u_3, \dots, u_m\}$ and $V(G_2) = \{v_1, v_2, v_3, \dots, v_n\}$. Now let $E_3 = \{u_i v_j : u_i \in V(G_1), v_j \in V(G_2)\}$. Let G_3 be the subgraph of $G_1 + G_2$ with the edge set E_3 . Therefore, $G_1 + G_2 = G_1 \cup G_2 \cup G_3$. Also, let f_1, f_2, f_3 be the IASIs defined on G_1, G_2, G_3 respectively. Given that f_1 and f_2 are strong IASIs. Let f be an IASI defined on $G_1 + G_2$ by

$f(v) = f_i(v)$ if $v \in V(G_i), i = 1, 2, 3$. Then, by Theorem 2.7, f is a strong IASI if and only if f_3 is a strong IASI.

First assume that f is a strong IASI. Then, f_3 is also a strong IASI. Hence, $|g_{f_3}(u_i v_j)| = |f_3(u_i)| \cdot |f_3(v_j)| \forall u_i \in V(G_1), v_j \in V(G_2)$. By Lemma 1.4, the difference sets of the set-labels of these vertices follow the relation $D_{u_i} < D_{v_j}, \forall u_i \in V(G_1), v_j \in V(G_2)$.

Conversely, assume that $D_{u_i} < D_{v_j}, \forall u_i \in V(G_1), v_j \in V(G_2)$. Then, $|g_{f_3}(u_i v_j)| = |f_3(u_i)| \cdot |f_3(v_j)| \forall u_i v_j \in E_3$. Therefore, f_3 is a strong IASI on G_3 and hence f is a strong IASI on $G_1 + G_2$. This completes the proof. $\square$

Theorem 2.11. *Let G_1 and G_2 be two strong IASI graphs. Then, $\kappa(G_1 + G_2) = \kappa(G_1) + \kappa(G_2)$.*

Proof. Let H_1 be a maximal clique with order m in G_1 and H_2 be a maximal clique with order n in G_2 . Since every vertex of G_1 is joined to every vertex of G_2 in $G_1 + G_2$, it is so in $H_1 + H_2$ also. Since H_1 and H_2 are cliques, they are complete graphs. Hence, every vertex of H_1 is adjacent to all other vertices of H_1 and all vertices of H_2 . Similarly, every vertex of H_2 is adjacent to all other vertices of H_2 and all vertices of H_1 . Hence, $H_1 + H_2$ is an $(m + n - 1)$ -regular graph on $m + n$ vertices. Therefore, $H_1 + H_2$ is a complete graph and hence is a clique in $G_1 + G_2$. Since H_1 and H_2 are maximal cliques, $H_1 + H_2$ is maximal in $G_1 + G_2$. Hence, $\kappa(G_1 + G_2) = m + n = \kappa(G_1) + \kappa(G_2)$. $\square$

Next, we discuss about the nourishing number of the complement of a strong IASI graph G . A graph G and its complement $\bar{G}$ have the same set of vertices and hence G and $\bar{G}$ have the same set-labels for their corresponding vertices. We observe that the strong IASIs, except some, defined on G do not induce strong IASI on $\bar{G}$. A set-labeling of $V(G)$ that defines a strong IASI for both the graphs G and its complement $\bar{G}$ may be called a *strongly concurrent set-labeling*. The set-labels of the vertex set of G mentioned in this section are strongly concurrent. Hence, we propose the following results.

Proposition 2.12. *Let G be a strong IASI graph on n vertices and let $\bar{G}$ be its complement. Then, $\bar{G}$ admits a strong IASI if and only if the length of the chain of difference sets of set-labels of vertices in G or in $\bar{G}$ is n .*

Proof. We have $G \cup \bar{G} = K_n$. Therefore, the length of the chain of difference sets in $G \cup \bar{G}$ is n . Since $\kappa(G \cup \bar{G}) \geq \max\{\kappa(G), \kappa(\bar{G})\}$, the length of the chain of difference sets of set-labels of vertices in G or in $\bar{G}$ must be n .

Conversely, assume that the length of the chain of difference sets of set-labels of vertices in G or in $\bar{G}$ is n . Then, length of the chain of difference sets in $G \cup \bar{G}$ is n , which is the maximum possible length of a chain of difference

sets. Therefore, both G and $\bar{G}$ admit strong IASI under the same set-labels for the vertices of G . This completes the proof. $\square$

Corollary 2.13. *If G is a self-complementary graph on n vertices, which admits a strong IASI, then $\kappa(G) = n$.*

Proof. Since G is self complementary, we have $G \cong \bar{G}$ and hence $\kappa(G) = \kappa(\bar{G})$. Therefore, $n = \kappa(G \cup \bar{G}) = \max\{\kappa(G), \kappa(\bar{G})\} = \kappa(G)$. That is, $\kappa(G) = \kappa(\bar{G}) = n$. Therefore, G and $\bar{G}$ admit strong IASI. This completes the proof. $\square$

2.3 Strong IASIs of Graph Products

In graph products, we may have to make certain number of copies of the graphs and to make suitable of attachments between these copies and the given graphs. Therefore, we have to establish a suitable IASI, if exists, to each of these copies. Hence, we make the following remark.

Remark 2.14. Let G_i is a copy of a given graph G , which appears in a graph product. Let $n.A = \{na_i : a_i \in A\}$, for $n \in \mathbb{N}_0$. Note that $n.A \neq nA$. If f is a strong IASI on G , then the i -th copy of G denoted by G_i has the set-label f_i where $f_i(v_i) = r.f(v)$, $r \in \mathbb{N}$, where v_i is the vertex in G_i corresponding to the vertex v in G . We observe that if two sets A and B are disjoint, then $n.A$ and $n.B$ are also disjoint. Hence, if f is a strong IASI of G , then f_i is a strong IASI of G_i .

In this section, we verify the admissibility of strong IASI by the cartesian product of two graphs. By the term *product of graphs* we mean the cartesian product of graphs. we recall the definition the cartesian product of two graphs as follows.

Let $G_1(V_1, E_1)$ and $G_2(V_2, E_2)$ be two graphs. Then, the *cartesian product* or simply *product* of G_1 and G_2 , denoted by $G_1 \square G_2$, is the graph with vertex set $V_1 \times V_2$ defined as follows. Let $u = (u_1, u_2)$ and $v = (v_1, v_2)$ be two points in $V_1 \times V_2$. Then, u and v are adjacent in $G_1 \square G_2$ whenever $[u_1 = v_1 \text{ and } u_2 \text{ is adjacent to } v_2]$ or $[u_2 = v_2 \text{ and } u_1 \text{ is adjacent to } v_1]$. If $|V_i| = p_i$ and $E_i = q_i$ for $i = 1, 2$, then $|V(G_1 \square G_2)| = p_1 p_2$ and $|E(G_1 \square G_2)| = p_1 q_2 + p_2 q_1$.

Remark 2.15. We observe that the product $G_1 \square G_2$ is obtained as follows. Make p_2 copies of G_1 . Denote these copies by $G_{1_i}, 1 \leq i \leq p_2$, which corresponds to the vertex v_i of G_2 . Now, join the corresponding vertices of two copies G_{1_i} and G_{1_j} if the corresponding vertices v_i and v_j are adjacent in G_2 . Thus, we view the product $G_1 \square G_2$ as a union of p_2 copies of G_1 and

a finite number of edges connecting two copies G_{1_i} and G_{1_j} of G_1 according to the adjacency of the corresponding vertices v_i and v_j in G_2 , where $1 \leq i, j \leq p_2, i \neq j$.

Hence, we make the following inferences on the admissibility of strong IASI by the product of two strong IASI graphs.

Proposition 2.16. *Let G_1 and G_2 be two strong IASI graphs. Then, the product $G_1 \square G_2$ admits a strong IASI if and only if the set-labels of corresponding vertices different copies of G_1 which are adjacent in $G_1 \square G_2$ are disjoint.*

Proof. Assume that the graphs G_1 and G_2 admit strong IASIs, say f and g respectively and $G_1 \square G_2$ admits a strong IASI, say F . Since f is a strong IASI on G , then by Remark 2.14, the function f_i defined on the i -th copy G_{1_i} of G_1 , by $f_i(v_i) = r.f(v)$, for some positive integer r , is a strong IASI on G_{1_i} . If the corresponding vertices of two copies G_{1_i} and G_{1_j} are adjacent in $G_1 \square G_2$, then $|g_F(v_i v_j)| = |F(v_i)| \cdot |F(v_j)| = |f_i(v_i)| \cdot |f_j(v_j)|$, $\forall v_i \in G_{1_i}, v_j \in G_{1_j}$. Hence, $D_{v_i} < D_{v_j}$. That is $f_i(v_i)$ and $f_j(v_j)$ are disjoint.

Conversely, assume that the set-labels of corresponding vertices different copies of G_1 which are adjacent in $G_1 \square G_2$ are disjoint. Also, the function f_i defined on the i -th copy G_{1_i} of G_1 , by $f_i(v_i) = r.f(v)$, for some positive integer r , is a strong IASI on G_{1_i} , $1 \leq i \leq |V(G_2)|$. Hence, the difference sets of the set-labels of all the adjacent vertices in $G_1 \square G_2$ are disjoint. Therefore, $G_1 \square G_2$ admits a strong IASI. $\square$

Proposition 2.17. *Let G_1 and G_2 be two graphs which admit strong IASIs. Then, $\kappa(G_1 \square G_2) = \max\{\kappa(G_1), \kappa(G_2)\}$.*

Proof. Let H_1 and H_2 be the maximal clique in G_1 and G_2 respectively. Without loss of generality, let H_1 be greater than H_2 in terms of the number of vertices in them. For $1 \leq i \leq |V(G_2)|$, let H_{1_i} be the copy of H_1 in G_{1_i} and is the maximal clique in G_{1_i} . Now, observe that no vertex of another copy G_{1_j} is adjacent to all the vertices of H_{1_i} . Since, all cliques H_{1_i} are isomorphic, for $1 \leq i \leq |V(G_2)|$, H_{1_i} is the maximal clique in $G_1 \square G_2$. Hence, $\kappa(G_1 \square G_2) = |V(H_{1_i})| = |V(H_1)|$. Therefore, in general, $\kappa(G_1 \square G_2) = \max\{\kappa(G_1), \kappa(G_2)\}$. $\square$

Next, let us recall the definition of corona of two graphs. The *corona* of two graphs G_1 and G_2 , denoted by $G_1 \odot G_2$, is the graph obtained taking one copy of G_1 (which has p_1 vertices) and p_1 copies of G_2 and then joining the i -th point of G_1 to every point in the i -th copy of G_2 .

The following results establish the admissibility of strong IASI by the corona of two strong IASI graphs and its nourishing number.

Theorem 2.18. *Let $G_1(V_1, E_1)$ and $G_2(V_2, E_2)$ be two strong IASI graphs. Then, their corona $G_1 \odot G_2$ admits a strong IASI if and only if the difference set of the set-label of every vertex in G_1 is disjoint from the difference sets of the set-labels of all vertices of the corresponding copy of G_2 .*

Proof. Let f and g be the strong IASI on G_1 and G_2 respectively. Let $g_i = r.g, 1 \leq i \leq |V(G_1)|$, r being a positive integer, be an IASI defined on the i -th copy G_i of G . Then, by Remark 2.14, g_i is a strong IASI on G_i for all $i = 1, 2, 3, \dots, |V(G_1)|$.

First assume that $G_1 \odot G_2$ admits a strong IASI. Also, each copy of G_2 admits strong IASIs. Since $G_1 \odot G_2$ admits a strong IASI, the difference set of the set-label of each vertex u_i of G_1 and the difference set of the set-label of each vertex v_{ji} of G_{2i} , where $1 \leq i \leq |V(G_1)|, 1 \leq j \leq |V(G_2)|$, must be disjoint.

Conversely, assume that the difference set of the set-label of every vertex in G_1 is disjoint from the difference sets of the set-labels of all vertices of the corresponding copy of G_2 . Since each copy of G_2 is also strong IASI graph, for every pair of adjacent vertices u, v in $G_1 \odot G_2$, the difference sets D_u and D_v hold the relation $D_u < D_v$. Hence, $G_1 \odot G_2$ admits a strong IASI. $\square$

Proposition 2.19. *If G_1 and G_2 are two strong IASI graphs, then*

$$\kappa(G_1 \odot G_2) = \begin{cases} \kappa(G_1) & \text{if } \kappa(G_1) > \kappa(G_2) \\ \kappa(G_2) + 1 & \text{if } \kappa(G_2) > \kappa(G_1) \end{cases}$$

Proof. Let H_1 and H_2 be the maximal cliques in G_1 and G_2 respectively. Then, H_{2i} is the copy of H_2 in G_{2i} , which is maximal in G_{2i} . Since the vertex u_i of H_1 is adjacent to all vertices of the copy H_{2i} in $G_1 \odot G_2$, we can find $|V(G_1)|$ cliques in $G_1 \odot G_2$ with clique number $1 + |V(H_{2i})| = 1 + |V(H_2)| = 1 + \kappa(G_2)$.

If $\kappa(G_1) > \kappa(G_2)$, then clearly, $\kappa(G_1 \odot G_2) = \kappa(G_1)$. If $\kappa(G_1) < \kappa(G_2)$, then the maximal clique in $G_1 \odot G_2$ is $H_{2i} + \{u_i\}$. Therefore, $\kappa(G_1 \odot G_2) = 1 + \kappa(G_2)$. This completes the proof. $\square$

3 Conclusion

In this paper, we have discussed about the admissibility of strong IASIs by certain graph classes, graph operations and graph products. We have established some results on the nourishing number of the graphs and graph

operations. The admissibility of strong IASI by various other graph operations and graph products and finding the corresponding nourishing numbers are remained to be estimated.

More properties and characteristics of strong IASIs, both uniform and non-uniform, are yet to be investigated. The problems of establishing the necessary and sufficient conditions for various graphs and graph classes to have certain IASIs still remain unsettled. All these facts highlight a great scope for further studies in this area.

References

- [1] R Aharoni, E Berger, M Chudnovsky and J Ziani, *Cliques in the Union of Graphs*, <http://www.columbia.edu/mc2775/edgeunion.pdf>
- [2] J A Bondy and U S R Murty, (1976). **Graph Theory with Applications**, North-Holland, New York.
- [3] R Frucht and F Harary (1970). *On the Corona of Two Graphs*, Aequationes Math., **4**(3), 322-325.
- [4] K A Germina and T M K Anandavally, (2012). *Integer Additive Set-Indexers of a Graph: Sum Square Graphs*, Journal of Combinatorics, Information and System Sciences, **37**(2-4), 345-358.
- [5] K A Germina, N K Sudev, (2013). *On Weakly Uniform Set-Indexers of Graphs*, Int.Math. Forum, **8**(37), 1827-34.
- [6] R Hammack, W Imrich and S Klavzar (2011). *Handbook of Product graphs*, CRC Press.
- [7] F Harary, **Graph Theory**, Addison-Wesley Publishing Company Inc., (1969).
- [8] W Imrich, S Klavzar, (2000). **Product Graphs: Structure and Recognition**, Wiley.
- [9] N K Sudev and K A Germina, *Some New results on Strong Integer Additive Set-indexers*, communicated.
- [10] D B West, (2001). **Introduction to Graph Theory**, Pearson Education Inc.